

Reinterpretation of gizzard sizes of red knots world-wide emphasises overriding importance of prey quality at migratory stopover sites

Jan A. van Gils^{1,2,*†}, Phil F. Battley^{1,‡}, Theunis Piersma^{1,2} and Rudi Drent²

¹Department of Marine Ecology and Evolution, Royal Netherlands Institute for Sea Research (NIOZ),
PO Box 59, 1790 AB Den Burg, Texel, The Netherlands

²Animal Ecology Group, Centre for Ecological and Evolutionary Studies (CEES), University of Groningen,
PO Box 14, 9750 AA Haren, The Netherlands

The size of digestive organs can be rapidly and reversibly adjusted to ecological circumstances, but such phenotypic flexibility comes at a cost. Here, we test how the gizzard mass of a long-distance migrant, the red knot (*Calidris canutus*), is adjusted to (i) local climate, (ii) prey quality and (iii) migratory fuelling demands. For eight sites around the world (both wintering and stopover sites), we assembled data on gizzard masses of free-living red knots, the quality of their prey and the local climate. Using an energetic cost–benefit approach, we predicted the gizzard size required for fastest fuelling (net rate-maximization, i.e. expected during migration) and the gizzard size required to balance daily energy budgets (satisficing, expected in wintering birds) at each site. The measured gizzards matched the net rate-maximizing predictions at stopover sites and the satisficing predictions at wintering sites. To our surprise, owing to the fact that red knots selected stopover sites with prey of particularly high quality, gizzard sizes at stopovers and at wintering sites were nevertheless similar. To quantify the benefit of minimizing size changes in the gizzard, we constructed a model incorporating the size-dependent energy costs of maintaining and carrying a gizzard. The model showed that by selecting stopovers containing high-quality prey, metabolic rates are kept at a minimum, potentially reducing the spring migratory period by a full week. By inference, red knots appear to time their stopovers so that they hit local peaks in prey quality, which occur during the reproductive seasons of the intertidal benthic invertebrates.

Keywords: benthos; *Calidris canutus*; digestive constraint; fuelling; gizzard; migration

1. INTRODUCTION

The realization that size, structure and function of digestive organs can be fine-tuned rapidly and reversibly to local/temporal ecological circumstances (phenotypic flexibility; reviewed by Piersma & Lindström 1997 and Piersma & Drent 2003), raises the question of costs that such adjustments entail. For example, as a response to increased energy demands through cold exposure, house wrens (*Troglodytes aedon*) increased the length of their small intestines (Dykstra & Karasov 1992). Quail (*Coturnix japonica*) increased gizzard size, intestine length and mucosal surface when food quality was reduced (Starck & Rahmaan 2003; and see López-Calleja & Bozinovic 2003 for a similar effect in hummingbirds, *Sephanoides sephanoides*). When fuelling for migration, garden warblers (*Sylvia borin*) increased their digestive tract size (Hume & Biebach 1996).

Long-distance migrants encounter a variety of climates and food qualities throughout the year and have different energy demands during non-breeding and migration seasons. These factors result in seasonal changes in food requirements (e.g. Piersma 2002). Given the existing empirical relationships between food processing rate and

digestive organ size (Van Gils *et al.* 2003), migrants provide an excellent model for studying how organ size is optimized in relation to external and internal energy demands.

It is increasingly realized that much of the reproductive success of long-distance migrant birds may depend on the ecological conditions encountered long before arrival on the breeding grounds (Ebbinge & Spaans 1995; Drent *et al.* 2003; Baker *et al.* 2004). In a world where many stopover sites are under threat, this gives the study of the selection pressures during migration more than academic interest. Arguments for the safeguarding of stopover sites usually come from observations of large concentrations (Wetlands International 2002), their provision of ample food (Myers *et al.* 1987) or theoretical arguments based on issues of speed of migration (Alerstam & Hedenström 1998). Here we combine these factors in a single argument based on the organ architecture of a shorebird species during different times of the year in relation to diet quality, energy expenditure and migratory phase.

We examine gizzard flexibility in the red knot (*Calidris canutus*), a medium-sized shorebird that undertakes several non-stop long-distance flights (1 400–6 500 km) between high-arctic breeding grounds and temperate, or tropical non-breeding grounds (Piersma *et al.* 2005). At these intertidal wintering sites, red knots feed primarily on hard-shelled prey that are generally of poor quality (i.e. low ratio of digestible to indigestible matter). Over the past decade, studies on red knots have quantified (i) the

* Author for correspondence (j.vangils@nioo.knaw.nl).

† Present address: Plant-Animal Interactions, Netherlands Institute of Ecology (NIOO-KNAW), Centre for Limnology, Rijksstraatweg 6, 3631 AC Nieuwersluis, The Netherlands.

‡ Present address: Department of Mathematics and Statistics, University of Otago, PO Box 56, Dunedin, New Zealand.




Figure 1. A map of the world with the knot's flyways (arrows), breeding grounds (dark grey surfaces) and wintering grounds (dots, scaled to estimated wintering population sizes). Distances (km) travelled between main wintering and breeding grounds are 9 000 (*canutus*), 4 680 (*islandica*), 15 000 (*rufa*), 15 000 (*rogersi*) and 10 400 (*piersmai*).

energetic costs of living in (Wiersma & Piersma 1994; Bruinzeel & Piersma 1998) and travelling between (Kvist *et al.* 2001) sites with contrasting climates (reviewed by Piersma 2002), (ii) the energetic costs and benefits of feeding on prey of different quality (Piersma *et al.* 2003; Van Gils *et al.* 2003, 2005a,b) and (iii) fuelling rates at different sites around the world (Gudmundsson *et al.* 1991; González *et al.* 1996; Piersma *et al.* 2005). Recent advances in a non-invasive technique (ultrasonography; see Dietz *et al.* 1999) enabled us to reveal flexibility in the size of the gizzard enforced by experimental changes in prey quality (Dekinga *et al.* 2001; Van Gils *et al.* 2003). This organ is vital in the feeding ecology of the knot as the hard-shelled prey, which provide most of their diet, are ingested whole and crushed in the muscular gizzard (Piersma *et al.* 1993b; Battley & Piersma 2005). Changes in gizzard size, which can occur rapidly and reversibly (50% within a week; Dekinga *et al.* 2001), are likely to have an impact on the knot's energy budget (Piersma *et al.* 2003; Van Gils *et al.* 2003, 2005b). As rates of shell crushing increase with gizzard size, energetic benefits increase with gizzard size (Van Gils *et al.* 2003). However, as larger gizzards require larger maintenance and transport costs, energetic costs should also increase with gizzard size (Piersma *et al.* 2003; Van Gils *et al.* 2003).


In this paper we predict optimal gizzard sizes at different sites based on energy demand and prey quality. Energy demand varies with (i) local climate and (ii) 'internal' energy demands for migratory fuelling. Prey quality affects the amount of bulk material that must be processed to meet the daily energy demands. Optimal gizzard size depends on whether birds are balancing gross energy intake and expenditure ('satisficing'; Nonacs & Dill 1993; expected during winter when there is no change in body mass), or maximizing net energetic benefits (gross intake minus expenditure; 'net rate-maximization'; Stephens & Krebs 1986; expected during spring when rapidly fuelling for migration). We tested this for five of the six recognized subspecies of knot (Piersma & Davidson 1992; Tomkovich 2001), for which data on climate, prey quality and gizzard size are available. These are *canutus* breeding at Taymyr Peninsula and wintering in west and southwest

Africa, *islandica* breeding in north Greenland and northeast Canada and wintering in northwest Europe, *rufa* breeding in the central Canadian arctic and wintering in southern Patagonia and Tierra del Fuego, *rogersi* breeding at Chukotskiy Peninsula and wintering in New Zealand and southeast Australia and *piersmai* breeding on the New Siberia islands and wintering in northwest Australia (figure 1). We considered shellfish-eaters only, but we will discuss exceptions later (horseshoecrab-egg-eating *rufa* knots during stopover in Delaware Bay; see Castro & Myers 1993). The links not specifically covered in this paper, owing to lack of data, are the movements of *roselaari* and the 6 900-km journey linking the South Africa *canutus* wintering sites with West African sites.

2. MATERIAL AND METHODS

(a) *Modelling cost-benefit as a function of gizzard size*
We modelled two types of gizzards. (i) The so-called 'satisficing' gizzards *balance* gross energy income with energy expenditure on a daily basis. (ii) The so-called 'net-rate maximizing' gizzards *maximize* net energy income on a daily basis (gross energy income minus energy expenditure). Of what size these two types of gizzard turn out to be, depends on how energetic benefits and energetic costs scale with gizzard size G . These relations are clarified in figure 2a and parameterized below.

We emphasize that we have followed the lead of optimal foraging theory (Stephens & Krebs 1986) by explicitly modelling *energetic* currencies in order to understand variation in gizzard mass. We realize that this somewhat limited approach ignores associated non-energetic costs, for example risk of predation, which may increase with gizzard size due to mass-dependent predation risk (MacLeod *et al.* 2005). Moreover, we neglect additional mortality costs that generally come with increased rates of energy assimilation (Yearsley *et al.* 2002) or expenditure (Daan *et al.* 1996), for example through an increased release of free radicals (Finkel & Holbrook 2000). We simply assume that birds either balance or maximize their energy budget, no matter the amount of energy this takes (an energetic currency that could partly account for expenditure-induced costs would be (the modified form of) efficiency, which expresses gross energy gained per unit of energy



expended (Houston 1995; Hedenström & Alerstam 1995)). The reasons that, in spite of these notions, we have focused on two *energetic* currencies only are: (i) simplicity and, most importantly, (ii) knowledge about parameter-values. In contrast to the body of knowledge that exists on energetic cost functions in knots (reviewed in §1), we have no clue how non-energetic mortality costs scale with gizzard size, although studies on predation-related cost functions in knots are being made (Van den Hout, in preparation). Thus, in order to make tractable and reliable predictions, we restricted ourselves to a cost-benefit model based on energetics only.

When modelling energetic *benefits* as a function of G we relied on the observation that the amount of shell material that can be crushed and processed per unit time increases quadratically with gizzard mass ($10^{-4.293}$ g shell mass $s^{-1} g^{-2}$ gizzard mass; Van Gils *et al.* 2003). Rates of metabolizable energy intake (MEIR; W) can therefore be lifted by increasing gizzard size (G ; g) and/or selecting prey with high flesh-to-shell ratios (prey quality Q ; J metabolizable energy g^{-1} shell mass). This can be formalized to $MEIR = 10^{-4.293} G^2 Q$ (Van Gils *et al.* 2003). See figure 2b for effects of prey quality on optimal gizzard size.

Prey quality Q was determined as precisely as possible by selecting only prey species and sizes fed upon by knots (see electronic supplementary material for diet composition and table 1 for resulting values of Q). Since knots ingest their shelled, largely indigestible prey whole, their faeces reveal much about their diet. Shell fragments can usually be identified to species-level and hinge-sizes allow reconstructions of prey size (Dekinga & Piersma 1993). In this way, hard parts of occasionally ingested crustaceans can also be found and identified (Van Gils *et al.* 2003, 2005b). Following this methodology, we reconstructed digestive prey quality for each subspecies at most sites, except for three where Q was extracted from the literature (*islandica* at the wintering site from Van Gils *et al.* (2003) and at stopover from Alerstam *et al.* (1992) and *canutus* and *rufa* at stopover from Van Gils *et al.* (2003) and González *et al.* (1996), respectively). Flesh and shell mass for each bivalve prey species and where possible each relevant size class, were determined. This was done by removing the flesh from the shells, drying both flesh and shell for three days to constant mass at 60°C and finally measuring dry masses to the nearest 0.1 mg (yielding DM_{flesh} and DM_{shell} , respectively). Next, the dried flesh was incinerated for 2 h at 550°C to determine ash-free dry mass ($AFDM_{\text{flesh}}$). After averaging $AFDM_{\text{flesh}}$ and DM_{shell} , we calculated Q as $daAFDM_{\text{flesh}}/DM_{\text{shell}}$, where d is energetic density of the flesh ($22 \text{ kJ g}^{-1} AFDM_{\text{flesh}}$; Zwartz & Wanink 1993) and a is assimilation efficiency (0.8; Kersten & Piersma 1987; Piersma 1994). Note that Piersma (1994) found that assimilation efficiencies did not vary between individual birds or between prey species.

In order to model energetic *costs* as a function of G , we used the linear relation between basal metabolic rate BMR (W) and lean body mass L (g) as observed by Piersma *et al.* (1996; $BMR = 0.0081 L - 0.046$) and assumed that L equals $100 \text{ g} + G +$ intestine mass (Van der Meer & Piersma 1994) and that gizzard mass scales to intestine mass in a one-to-one isomorphic relationship (as observed by Piersma *et al.* 2003), or formally $L = 100 + 2G$. The metabolic rate due to flying MR_{fly} (W) scales to body mass B (g) as $MR_{\text{fly}} = 10^{0.39} B^{0.35} - 0.95$ (Kvist *et al.* 2001). Metabolic rate due to walking MR_{walk} (W) scales to body mass B (g) in the following manner: $MR_{\text{walk}} = v840(B/100)^{2.90}$ (Bruinzeel *et al.* 1999), where v is a walking speed of 0.072 m s^{-1} (Piersma *et al.* 2003). In order

Table 1. Site and sample data for the five subspecies of red knot, with predicted TRCs ($W; \pm s.e.$), average digestive prey quality ($\text{kJ g}^{-1} \text{DM}_{\text{shell}} \pm s.e.$), expected satisficing and net rate-maximizing gizzard masses ($g; s.e.$ -confidence interval in brackets), the number of gizzards sampled and the observed gizzard masses ($g; \text{mean} \pm s.e.$). (References are to detailed studies at the particular site.)

| subspecies | migratory phase | site | month | TRC (W) | prey quality ($\text{kJ g}^{-1} \text{DM}_{\text{shell}}$) | exp. satisficing gizzard mass (g) | exp. net rate-maximizing gizzard mass (g) | N gizzards sampled | obs. mean gizzard mass $\pm s.e.$ (g) | reference |
|------------------|-----------------|--------------------------------|---------------|-----------------|--|---------------------------------------|---|----------------------|---|---|
| <i>canutus</i> | wintering | Banc d'Arguin, Mauritania | December | 1.49 ± 0.27 | 0.89 ± 0.07 | $10.51 (9.70-11.38)$ | $16.64 (15.71-17.67)$ | 6 | 9.89 ± 0.53 | this study |
| | stopover | Wadden Sea, NW-Europe | May | 1.99 ± 0.03 | 3.74 ± 0.03 | $5.73 (5.68-5.78)$ | $8.14 (7.93-8.34)$ | 2 | 8.00 ± 0.30 | Van Gils <i>et al.</i> (2003) |
| <i>islandica</i> | wintering | Wadden Sea, NW-Europe | January | 2.93 ± 0.02 | 2.10 ± 0.03 | $8.49 (8.40-8.58)$ | $10.86 (10.55-11.17)$ | 60 | 8.75 ± 0.18 | Van Gils <i>et al.</i> (2003) |
| <i>islandica</i> | stopover | Selvogur, SW-Iceland | May | 2.26 ± 0.04 | 2.35 ± 0.31 | $6.33 (5.91-6.85)$ | $9.45 (8.70-10.35)$ | 8 | 7.95 ± 0.48 | Piersma <i>et al.</i> (1999), Alerstam <i>et al.</i> (1992) |
| <i>rufa</i> | wintering | Tierra del Fuego, S-Argentina | February | 1.82 ± 0.06 | 2.03 ± 0.14 | $7.21 (6.92-7.53)$ | $11.04 (10.47-11.66)$ | 13 | 8.09 ± 0.20 | this study |
| | stopover | San Antonio Oeste, E-Argentina | March | 1.35 ± 0.05 | 3.50 ± 0.23 | $5.09 (4.89-5.31)$ | $8.41 (7.97-8.87)$ | 7 | 8.17 ± 0.20 | González <i>et al.</i> (1996) |
| <i>rogersi</i> | wintering | Northern New Zealand | March | 1.30 ± 0.03 | 0.82 ± 0.04 | $9.55 (9.29-9.84)$ | $15.49 (14.82-16.20)$ | 5 | 9.31 ± 0.64 | Battley & Piersma (1997) |
| <i>piersmai</i> | wintering | Roebuck Bay, NW-Australia | January-April | 0.67 ± 0.04 | 2.22 ± 0.30 | $5.91 (5.53-6.39)$ | $10.55 (9.70-11.59)$ | 24 | 5.94 ± 0.36 | this study |

to equate these mass-dependent equations as a function of gizzard mass, we assume that $B=L$ (i.e. no fat is assumed; as fat is metabolically inactive this assumption hardly affects the calculations through the relatively small transport costs). Metabolic rate owing to probing equals 0.47 W (Piersma *et al.* 2003) and heat increment of feeding (HIF) amounts to 5.2 kJ g^{-1} AFDM_{flesh} digested (Piersma *et al.* 2003).

Finally, thermoregulatory cost (TRC) is estimated on the basis of Wiersma & Piersma (1994), who provide (i) direct estimates of TRC for *islandica* and *canutus* and (ii) an estimate of thermal conductance that we used to translate ambient temperatures into TRC for the other three subspecies (listed in table 1; using data in Stans *et al.* 1972). This aspect of cost is exemplified in figure 2c. We assume that some heat-generating sources can contribute towards TRC, which makes life in the cold considerably cheaper. Walking generates most of its heat close to the skin and therefore only 30% of it can be 'kept' and used as a substitute for TRC (Bruinzeel & Piersma 1998). Since HIF represents heat generated in the core of the body, 100% of it can potentially substitute for TRC (and does, K. Jalvingh & F. Vézina unpublished data). The 'maintenance cost' as mostly used by Wiersma & Piersma (1994) and Piersma (2002) is embodied in the well known 'Scholander-curve' (Scholander *et al.* 1950) which equals TRC below and BMR above the lower critical temperature (Scholander *et al.* 1950).

To complete the calculations on the *daily* energy budget, we multiplied these expenditure and intake rates with daily time allocations to foraging, flying and resting. Red knots feed at exposed intertidal mudflats during day and night and therefore have daily foraging periods of 12 h on average (Piersma *et al.* 1994; Van Gils & Piersma 1999; Van Gils *et al.* 2000). This value was used for all subspecies at all sites except for *islandica* stopping over at Iceland (14.2 h; Alerstam *et al.* 1992) and for *rogersi*




Figure 3. Daily energy requirement (horizontal axis) and prey quality (diagonal lines) predict the daily mass of shell material processed (left vertical axis) and thus predict the required gizzard size (right vertical axis). (a) The prediction for satisficing and rate-maximizing *C. c. canutus*, both at wintering (filled squares) and stopover sites (open squares). Arrows indicate the expected change in gizzard mass when changing from wintering/satisficing conditions to wintering/rate-maximizing or to stopover/rate-maximizing conditions. (b) Predictions for all subspecies (ignoring rate-maximization during winter). Due to an increase in prey quality, little change in gizzard size is required when shifting from wintering to stopover sites (arrows). Note that the gizzard scale is based on 12 h of foraging per day, which holds for all cases except for wintering *rogersi* (15 h) and stopping-over *islandica* (14.2 h). In these two cases, predicted gizzard mass is actually somewhat smaller than plotted here ($\sqrt{12/15}$ and $\sqrt{12/14.2}$ times the plotted gizzard mass).

largest in wintering *canutus* (figure 4; table 1). Gizzards at wintering sites did not differ from predicted satisficing sizes ($p > 0.15$, using the 'combined probability test' as proposed by Sokal & Rohlf (1995); we compared test statistic $-2\sum \ln p = 13.74$ with $\chi^2_{2 \times 5}$) and were smaller than the predicted net rate-maximizing sizes ($p < 0.0005$, i.e. comparing $-2\sum \ln p = 63.58$ with $\chi^2_{2 \times 3}$; figure 4; table 1). Conversely, gizzards at stopover sites did not differ from predicted net rate-maximizing sizes ($p > 0.4$, i.e. comparing $-2\sum \ln p = 5.90$ with $\chi^2_{2 \times 3}$) and were larger than the predicted satisficing sizes ($p < 0.0005$, i.e. comparing $-2\sum \ln p = 129.03$ with $\chi^2_{2 \times 5}$; figure 4; table 1). However, stopover gizzards were of similar size to the wintering gizzards ($p > 0.1$, taking differences between subspecies into account, $N = 125$, $R^2 = 0.41$).

4. DISCUSSION

Gizzards at stopovers and gizzards at wintering sites were of similar size (figure 4). This may seem counterintuitive, especially since 'stopover gizzards' were of rate-maximizing




Figure 4. Observed gizzard masses (mean \pm s.e.) for each subspecies at its wintering site (closed symbol) and stopover site (open symbol, where data are available). Boxes in the background give range in expected gizzard mass (\pm s.e.), where each box spans the expected gizzard mass of satisficing (bottom) and net rate-maximizing birds (top).

size while 'wintering gizzards' were of satisficing size (figure 4; table 1). However, prey qualities at stopovers were on average twice those at wintering sites (figure 3b; $N = 8$; $p < 0.05$). This enabled gizzards of the same size to accommodate a maximal fuelling mode (arrows in figure 3b). When migrating northwards, the increase in prey quality for *canutus* is of such magnitude (from 0.89 to 3.74 $\text{kJ g}^{-1} \text{DM}_{\text{shell}}$) that fuelling 'spring gizzards' are predicted and observed to be 2 g smaller than satisficing 'winter gizzards'. With an increase in prey quality from 2.03 to 3.50 $\text{kJ g}^{-1} \text{DM}_{\text{shell}}$, it seems as if *rufa* uses its relatively nearby stopover in San Antonio Oeste ('only' 1400 km away from the wintering grounds at Tierra del Fuego) as a springboard to escape from the Patagonian wintering grounds. The increase in prey quality is smallest in the temperate-zone wintering *islandica* (from 2.10 to 2.35 $\text{kJ g}^{-1} \text{DM}_{\text{shell}}$), but in this case fuelling in Iceland with relatively small gizzards seems feasible by foraging longer than 12 h (Alerstam *et al.* 1992), although not at maximal rates (see figure 4).

Apparently, red knots only utilize the stopover sites that harbour prey of high quality. This agrees with theoretical predictions (Alerstam & Lindström 1990) and other empirical studies on red knots (those listed by Gudmundsson *et al.* 1991; Van Gils *et al.* 2005b): poor-quality stopovers ought to be skipped and indeed are. In an attempt to quantify the benefits of selecting high-quality stopovers, we modelled total duration of migration as a function of stopover selection for a (theoretical) knot migrating 15 000 km (figure 5). The model builds on our previous calculations, but now includes fattening. Fuelling birds deposit up to 100 g of fat at a biosynthesis-cost of 0.33 J per J of tissue (Ricklefs 1974). The next step in our model is their long-distance flight. During long-distance flight, the burning of 1 g of fat yields 40 kJ. To take into account the instantaneous changes in energetic costs with changing amounts of fat (increasing when fuelling and decreasing when flying), the model runs at discrete time steps of 1 s. Across any prey quality, net-rate maximizing gizzards always yield a gross rate of energy gain that equals




Figure 5. The predicted effect of digestive prey quality at stopovers on overall speed for a theoretical, but realistic, knot population to migrate 15 000 km from wintering to breeding grounds. (a) Expected net-rate maximizing gizzard mass declines with prey quality (solid curved line; see eq. given in §2). Given the observed frequency distribution of prey qualities among sites (bars on horizontal axis), knots only select stopovers that harbour the highest prey qualities (light grey bars; dark grey bars represent wintering sites). As predicted, gizzards observed at stopovers are distributed around 8 g (bars on vertical axis). Mean stopover gizzard mass versus mean stopover prey quality is given by open dot (bars are 95% confidence intervals). (b) Metabolic rates are predicted to increase as a function of gizzard size, especially during flight. (c) Through these higher energetic costs, the total length of the migratory period increases with increasing gizzard size (i.e. declines with increasing prey quality). Grey band across the three figures shows that, according to the model, stopping over in the best instead of in the worst sites should reduce length of migratory period by about a week. Downsizing the gizzard before flight (to 6 g with no time or energy costs of transformation), results in only a minor reduction of the migratory period (dashed line in *b* and *c*).

the upper physiological limit as empirically derived by Kirkwood (1983) and Kvist & Lindström (2003; grey bars in figures 2 and 3). However, as net-rate maximizing gizzard size declines with increasing prey quality (figure 5a cf. figure 2b), the overall cost of transport and maintenance declines likewise with increasing prey quality (figure 5b).

This cost-reduction by minimizing gizzard size brings two advantages that should, according to the model, reduce overall duration of migration (by about one day per gram decline in gizzard mass; figure 5c). First, at the stopovers, higher *net* rates of energy gain (= gross gain minus costs) are achieved, leading to shorter stopover times (P_{dep} increases, using the terminology of Hedenström & Ålerstam 1998). Second, during flight, metabolic rate is reduced and thus longer distances can be travelled per gram of stored fat, which reduces required stopover time per distance travelled (P_{flight} decreases, using the terminology of Hedenström & Ålerstam 1998). Even though this second effect is stronger than the first on an instantaneous basis (given by differences in slopes in figure 5b), overall, most energy is saved at stopovers, since much more time and thus energy, is spent at stopovers than in flight (cf. Wikelski *et al.* 2003). Summarizing, overall speed of migration should improve by selecting stopovers that harbour high-quality prey. The overall speed of migration expected by the model (15 000 km in 91 days = 165 km day⁻¹) approaches the value of 175 km day⁻¹ actually observed in red knots (Hedenström & Ålerstam 1998).

We extended the model by including the flexibility to further reduce gizzard size just before the onset of a long-distance flight down to 6 g (as observed by Battley & Piersma 1997; Piersma *et al.* 1999). We assumed such transformations incur no time or energy costs. Recent evidence suggests that the energetic costs involved may indeed be minor. Overgaard *et al.* (2002) found that growth of gastrointestinal organs in pythons is associated with vast increases in metabolic rate, but showed that such increases

are merely owing to the act of processing a meal rather than to the organ growth itself. Starck & Rahmaan (2003) found that (resting) metabolic rates in a bird species, quail, remain unaltered during size changes in gizzard and small intestine. Although energetic costs involved in size changes in both gizzard and intestine may thus be minute, Starck & Rahmaan additionally showed that the cellular processes underlying such changes differ substantially between both organs. Gizzards grow because of cell growth while intestines grow because of cell proliferation.

Due to the fact that fuelling gizzards are relatively small (8 g), the 25% downscaling to 6 g only marginally improves overall speed of migration through small reductions in P_{flight} (by only a single day over the full distance of 15 000 km, so the overall speed of migration = 167 km day⁻¹; dashed line in figure 5b,c). The far greater effect of prey quality remains.

Prey is of best quality (high amounts of flesh) in the reproductive phase just before the release of gametes (Gabbot 1983; Zwarts 1991; Honkoop & Van der Meer 1997). In temperate zones, reproduction of marine invertebrates is seasonally synchronized and takes place in spring (see references in Zwarts 1991), which is reflected in the peak in prey quality during spring (Van Gils *et al.* 2003). Keeping in mind that the boreal spring and thus prey reproduction starts later in the year with increasing latitude (Piersma *et al.* 1994), spring-migrating knots seem to follow a northwards 'wave' in prey quality. We therefore suggest that knots locate and time their stopovers to coincide with local peaks in prey reproductive periods. This idea has been suggested earlier with respect to increased availability of benthic prey (Piersma *et al.* 1994) and for Arctic-breeding waterfowl tuning their migration to plant growth ('the green wave hypothesis'; Drent *et al.* 1978).

In tropical regions, prey reproduction is much less synchronized and occurs throughout the year so that a distinct seasonal peak in prey quality may be absent (De Goeij *et al.* 2003). Therefore, rapid fuelling may be

impossible around the equator. This might explain why tropical regions are probably skipped as stopovers by knots wintering in the Southern Hemisphere (*rufa* and *rogersi*). Tropically wintering knots (*canutus* and *piersmai*) may find fuelling and leaving these sites difficult, a suggestion that has been made before (Ens *et al.* 1990; Piersma *et al.* 2005). Indeed, rates of fuelling are lower ($N = 14$ sites, $R^2 = 0.29$, $p < 0.05$) and length of fuelling periods are longer ($N = 14$ sites, $R^2 = 0.61$, $p = 0.001$) in tropical than in temperate regions (data from Piersma *et al.* 2005). West-African wintering *canutus* may face the severest problems. If their prey in spring had the same low quality observed in winter ($0.89 \text{ kJ g}^{-1} \text{ DM}_{\text{shell}}$), then the maximum fuelling rate would require a gizzard of approximately 17 g (figure 3a; table 1). In addition to higher maintenance and transport costs associated with such huge gizzards, 'space' for fat deposition might be limited in such muscular bodies. Indeed, knots departing from Banc d'Arguin are relatively light (average 168 g, $p < 0.05$ when comparing this with 13 other average departure masses given by Piersma *et al.* (2005)). Possibly, *canutus* manages with gizzards smaller than 17 g by extending the low tide period beyond the 'usual' 12 h day^{-1} (Zwarts *et al.* (1990); this has been observed for *islandica* in the Wadden Sea (Van Gils *et al.* 2005b) and Iceland (Alerstam *et al.* 1992)).

Fuelling in the Southern Hemisphere takes place during the austral autumn, a time of year long past the peak in prey quality in the (austral) spring (Wilson & Hodgkin 1967; Wilson 1969). Our 'prey-reproduction' hypothesis would therefore predict that among sites in temperate regions, fuelling and consequently speed of northward migration should proceed faster in the Northern Hemisphere than in the Southern Hemisphere. Using data published by Piersma *et al.* (2005), this prediction is upheld, both with respect to fuelling rates ($N = 11$ sites, $R^2 = 0.39$, $p < 0.05$) and with respect to lengths of fuelling periods ($N = 11$ sites, $R^2 = 0.36$, $p = 0.05$).

The approach used here, predicting optimal gizzard sizes from a prey's flesh-to-ballast ratio, is only applicable to knots that feed on hard-shelled mollusc prey. Applying the model to *rufa* feeding on the super-high quality eggs of horseshoe crabs (*Limulus polyphemus*) at its stopover in Delaware Bay (North America), predicted net-rate maximizing gizzards of about 1 g. The fact that observed gizzards were much larger than that (mean \pm s.e. = 7.0 ± 0.2 g, $N = 61$), hints that the grinding of horseshoe crab eggs is a fundamentally different process when compared to crushing the outer shells of molluscs. Only Delaware Bay red knots have small stones in their gizzards, presumably to grind the leathery surface of the eggs (T.P. unpubl. data; see Piersma *et al.* 1993b). Direct experimentation is called for to clarify this.

5. CONCLUSION

Variation in gizzard mass of wintering red knots is large (from 6 g in *piersmai* to almost 10 g in *canutus* and *rogersi*). The variation appears to reflect the best possible size adjustments to global variations in prey quality and climate in order to balance energy budgets on a daily basis. Variation in gizzard mass is, by contrast, small in migrating knots (around 8 g in all populations). By the selection of stopover sites with high-quality prey, red knots maximize fuelling rates and overall speed of migration

using relatively small gizzards. This idea generates testable predictions on the prey quality at as yet unexplored stopover sites, such as those of *rogersi* and *piersmai* along the East Asian–Australasian Flyway. Given the extremely late departure from their northwest Australian wintering site (Battley *et al.* 2005), *piersmai* is likely to encounter 'super-food' at its stopover in the Yellow Sea, enabling gizzards to be kept small and migration sped up.

This study is an outcome of many years of co-operative studies on red knots and we thank all involved. More specifically, we are very grateful to Graciela Escudero, Danny Rogers, Piet van den Hout, Casper Kraan, Anita Koolhaas and Maarten Brugge for making data available on prey quality, to Anne Dekinga and Maurine Dietz for expert ultrasonography, and to Jan Drent, Pieter Honkoop, Marcel Klaassen, Åke Lindström, Danny Rogers, Popko Wiersma, and two anonymous referees for comments on drafts. Dick Visser is acknowledged for the lay-out of the figures. This study was funded by a PIONIER-grant from the Netherlands Organisation for Scientific Research (NWO) to T.P.

REFERENCES

- Alerstam, T. & Hedenström, A. 1998 The development of bird migration theory. *J. Avian Biol.* **29**, 343–369.
- Alerstam, T. & Lindström, Å. 1990 Optimal bird migration: the relative importance of time, energy and safety. In *Bird migration: physiology and ecophysiology* (ed. E. Gwinner), pp. 331–351. Berlin: Springer-Verlag.
- Alerstam, T., Gudmundsson, G. A. & Johannesson, K. 1992 Resources for long distance migration: intertidal exploitation of *Littorina* and *Mytilus* by knots *Calidris canutus* in Iceland. *Oikos* **65**, 179–189.
- Baker, A. J., González, P. M., Piersma, T., Niles, L. J., De Lima Serrano do Nascimento, I., Atkinson, P. W., Clark, N. A., Minton, C. D. T., Peck, M. K. & Aarts, G. 2004 Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proc. R. Soc. B* **271**, 875–882. (doi:10.1098/rspb.2003.2663.)
- Battley, P. F. & Piersma, T. 1997 Body composition of Lesser Knots (*Calidris canutus rogersi*) preparing to take off on migration from northern New Zealand. *Notornis* **44**, 137–150.
- Battley, P. F. & Piersma, T. 2005 Adaptive interplay between feeding ecology and features of the digestive tract in birds. In *Physiological and ecological adaptations to feeding in vertebrates* (ed. J. M. Starck & T. Wang), pp. 201–228. Enfield, NH: Science Publishers.
- Battley, P. F., Rogers, D. I., Van Gils, J. A., Piersma, T., Hassell, C. J., Boyle, A & Yang, H. Y. 2005 How do red knots *Calidris canutus* leave Northwest Australia in May and reach the breeding grounds in June? Predictions of stopover times, fuelling rates and prey quality in the Yellow Sea. *J. Avian Biol.* **36**, 494–500. (doi:10.1111/j.2005.0908-8857.03730.x.)
- Bruinzeel, L. W. & Piersma, T. 1998 Cost reduction in the cold: heat generated by terrestrial locomotion partly substitutes for thermoregulation costs in Knot *Calidris canutus*. *Ibis* **140**, 323–328.
- Bruinzeel, L. W., Piersma, T. & Kersten, M. 1999 Low costs of terrestrial locomotion in waders. *Ardea* **87**, 199–205.
- Castro, G. & Myers, J. P. 1993 Shorebird predation on eggs of horseshoe crabs during spring stopover on Delaware Bay. *Auk* **110**, 927–930.
- Daan, S., Deerenberg, C. & Dijkstra, C. 1996 Increased daily work precipitates natural death in the kestrel. *J. Anim. Ecol.* **65**, 539–544.

- De Goeij, P., Lavaleye, M., Pearson, G. B. & Piersma, T. 2003 *Seasonal changes in the macro-zoobenthos of a tropical mudflat*. NIOZ-report 2003-4, Texel, The Netherlands.
- Dekinga, A. & Piersma, T. 1993 Reconstructing diet composition on the basis of faeces in a mollusc-eating wader, the knot *Calidris canutus*. *Bird Study* **40**, 144–156.
- Dekinga, A., Dietz, M. W., Koolhaas, A. & Piersma, T. 2001 Time course and reversibility of changes in the gizzards of red knots alternatively eating hard and soft food. *J. Exp. Biol.* **204**, 2167–2173.
- Dietz, M. W., Dekinga, A., Piersma, T. & Verhulst, S. 1999 Estimating organ size in small migrating shorebirds with ultrasonography: an intercalibration exercise. *Physiol. Biochem. Zool.* **72**, 28–37. (doi:10.1086/316648.)
- Drent, R., Ebbinge, B. & Weijand, B. 1978 Balancing the energy budgets of arctic-breeding geese throughout the annual cycle: a progress report. *Verh. Orn. Ges. Bayern* **23**, 239–264.
- Drent, R., Both, C., Green, M., Madsen, J. & Piersma, T. 2003 Payoffs and penalties of competing migratory schedules. *Oikos* **103**, 274–292. (doi:10.1034/j.1600-0706.2003.12274.x.)
- Dykstra, C. R. & Karasov, W. H. 1992 Changes in gut structure and function of house wrens (*Troglodytes aedon*) in response to increased energy demands. *Physiol. Zool.* **65**, 422–442.
- Ebbinge, B. S. & Spaans, B. 1995 The importance of body reserves accumulated in spring staging areas in the temperate zone for breeding in dark-bellied brent geese *Branta b. bernicla* in the high Arctic. *J. Avian Biol.* **26**, 105–113.
- Ens, B. J., Piersma, T., Wolff, W. J. & Zwarts, L. 1990 Homeward bound: problems waders face when migrating from the Banc d'Arguin, Mauritania, to their northern breeding grounds in spring. *Ardea* **78**, 1–7.
- Finkel, T. F. & Holbrook, N. J. 2000 Oxidants, oxidative stress and the biology of aging. *Nature* **408**, 239–247. (doi:10.1038/35041687.)
- Gabbot, P. A. 1983 Developmental and seasonal metabolic activities in marine molluscs. In *The mollusca. Vol. 2. Environmental biochemistry and physiology* (ed. P. W. Hochachka), pp. 165–217. New York, NY: Academic Press.
- González, P. M., Piersma, T. & Verkuil, Y. 1996 Food, feeding, and refuelling of red knots during northward migration at San Antonio Oeste, Rio Negro, Argentina. *J. Field Ornithol.* **67**, 575–591.
- Gudmundsson, G. A., Lindström, Å. & Alerstam, T. 1991 Optimal fat loads and long distance flights by migrating knots *Calidris canutus*, sanderlings *C. alba* and turnstones *Arenaria interpres*. *Ibis* **133**, 140–152.
- Hammond, K. A. & Diamond, J. 1997 Maximum sustained energy budgets in humans and animals. *Nature* **386**, 457–462. (doi:10.1038/386457a0.)
- Hedenström, A. & Alerstam, T. 1995 Optimal flight speeds of birds. *Phil. Trans. R. Soc. B* **348**, 471–487.
- Hedenström, A. & Alerstam, T. 1998 How fast can birds migrate? *J. Avian Biol.* **29**, 424–432.
- Honkoop, P. J. C. & Van der Meer, J. 1997 Reproductive output of *Macoma balthica* populations in relation to winter-temperature and intertidal-height mediated changes of body mass. *Mar. Ecol. Prog. Ser.* **149**, 155–162.
- Houston, A. I. 1995 Energetic constraints and foraging efficiency. *Behav. Ecol.* **6**, 393–396.
- Hume, I. D. & Biebach, H. 1996 Digestive tract function in the long-distance migratory garden warbler. *Sylvia borin*. *J. Comp. Physiol. B* **166**, 388–395. (doi:10.1007/s003600050024.)
- Karasov, W. H. & McWilliams, S. R. 2005 Digestive constraints in mammalian and avian ecology. In *Physiological and ecological adaptations to feeding in vertebrates* (ed. J. M. Starck & T. Wang), pp. 87–112. Enfield, NH: Science Publishers.
- Kersten, M. & Piersma, T. 1987 High levels of energy expenditure in shorebirds; metabolic adaptations to an energetically expensive way of life. *Ardea* **75**, 175–187.
- Kirkwood, J. K. 1983 A limit to metabolizable energy intake in mammals and birds. *Comp. Biochem. Physiol. A* **75**, 1–3. (doi:10.1016/0300-9629(83)90033-6.)
- Kvist, A. & Lindström, Å. 2003 Gluttony in migratory waders—unprecedented energy assimilation rates in vertebrates. *Oikos* **103**, 397–402. (doi:10.1034/j.1600-0706.2003.12259.x.)
- Kvist, A., Lindström, Å., Green, M., Piersma, T. & Visser, G. H. 2001 Carrying large fuel loads during sustained bird flight is cheaper than expected. *Nature* **413**, 730–731. (doi:10.1038/35099556.)
- López-Calleja, M. V. & Bozinovic, F. 2003 Dynamic energy and time budgets in hummingbirds: a study in *Sephanoides sephanioides*. *Comp. Biochem. Physiol. A* **134**, 283–295.
- MacLeod, R., Barnett, P., Clark, J. A. & Cresswell, W. 2005 Body mass change strategies in blackbirds *Turdus merula*: the starvation–predation risk trade-off. *J. Anim. Ecol.* **64**, 292–302. (doi:10.1111/j.1365-2656.2005.00923.x.)
- Myers, J. P., Morrison, R. I. G., Antas, P. T. Z., Harrington, B. A., Lovejoy, T. E., Sallaberry, M., Senner, S. E. & Tarak, A. 1987 Conservation strategy for migratory species. *Am. Sci.* **75**, 18–26.
- Nonacs, P. & Dill, L. M. 1993 Is satisficing an alternative to optimal foraging theory? *Oikos* **67**, 371–375.
- Overgaard, J., Andersen, J. B. & Wang, T. 2002 The effects of fasting duration on the metabolic response to feeding in *Python molurus*: an evaluation of the energetic costs associated with gastrointestinal growth and upregulation. *Physiol. Biochem. Zool.* **75**, 360–368. (doi:10.1086/342769.)
- Piersma, T. 1994 *Close to the edge: energetic bottlenecks and the evolution of migratory pathways in Knots*. Den Burg, Texel, The Netherlands: Uitgeverij Het Open Boek.
- Piersma, T. 2002 Energetic bottlenecks and other design constraints in avian annual cycles. *Integr. Comp. Biol.* **42**, 51–67.
- Piersma, T. & Davidson, N. C. 1992 The migrations and annual cycles of five subspecies of Knots in perspective. *Wader Study Group Bull.* **64**(Suppl.), 187–197.
- Piersma, T. & Drent, J. 2003 Phenotypic flexibility and the evolution of organismal design. *Trends Ecol. Evol.* **18**, 228–233. (doi:10.1016/S0169-5347(03)00036-3.)
- Piersma, T. & Lindström, Å. 1997 Rapid reversible changes in organ size as a component of adaptive behaviour. *Trends Ecol. Evol.* **12**, 134–138. (doi:10.1016/S0169-5347(97)01003-3.)
- Piersma, T., Hoekstra, R., Dekinga, A., Koolhaas, A., Wolf, P., Battley, P. & Wiersma, P. 1993a Scale and intensity of intertidal habitat use by knots *Calidris canutus* in the western Wadden Sea in relation to food, friends and foes. *Neth. J. Sea Res.* **31**, 331–357. (doi:10.1016/0077-7579(93)90052-T.)
- Piersma, T., Koolhaas, A. & Dekinga, A. 1993b Interactions between stomach structure and diet choice in shorebirds. *Auk* **110**, 552–564.
- Piersma, T., Verkuil, Y. & Tulp, I. 1994 Resources for long-distance migration of knots *Calidris canutus islandica* and *C. c. canutus*: how broad is the temporal exploitation window of benthic prey in the western and eastern Wadden Sea? *Oikos* **71**, 393–407.
- Piersma, T., Bruinzeel, L., Drent, R., Kersten, M. & Van der Meer, J. 1996 Variability in basal metabolic rate of a long-distance migrant shorebird (red knot, *Calidris canutus*) reflects shifts in organ sizes. *Physiol. Zool.* **69**, 191–217.

- Piersma, T., Gudmundsson, G. A. & Lilliendahl, K. 1999 Rapid changes in the size of different functional organ and muscle groups during refueling in a long-distance migrating shorebird. *Physiol. Biochem. Zool.* **72**, 405–415. (doi:10.1086/316680.)
- Piersma, T., Dekinga, A., Van Gils, J. A., Achterkamp, B. & Visser, G. H. 2003 Cost-benefit analysis of mollusc-eating in a shorebird. I. Foraging and processing costs estimated by the doubly labelled water method. *J. Exp. Biol.* **206**, 3361–3368. (doi:10.1242/jeb.00545.)
- Piersma, T., Rogers, D. I., González, P. M., Zwarts, L., Niles, L. J., De Lima Serrano Donascimento, I., Minton, C. D. T. & Baker, A. J. 2005 Fuel storage rates before northward flights in Red Knots world-wide: facing the severest ecological constraint in tropical intertidal environments. In *Birds of two worlds* (ed. R. Greenberg & P. P. Marra), pp. 262–273. Baltimore, MD: Johns Hopkins University Press.
- Ricklefs, R. E. 1974 energetics of reproduction in birds. In *Avian energetics* (ed. R. A. Paynter), pp. 152–292. Cambridge, MA: Nuttall Ornithological Club.
- Rogers, D. I. 2003 High-tide roost choice by coastal waders. *Wader Study Group Bull.* **100**, 73–79.
- Scholander, P. F., Hock, R., Walters, V. & Irving, L. 1950 Adaptation to cold in arctic and tropical mammals and birds in relation to body temperature, insulation, and basal metabolic rate. *Biol. Bull.* **99**, 259–271.
- Sokal, R. R. & Rohlf, F. J. 1995 *Biometry*, 3rd edn. New York, NY: W.H. Freeman.
- Stans, M. H., White, R. M. & Jacobs, W. C. 1972 *Climates of the world*. Washington, DC: U.S. Government Printing Office.
- Starck, J. M. & Rahmaan, G. H. A. 2003 Phenotypic flexibility of structure and function of the digestive system of Japanese quail. *J. Exp. Biol.* **206**, 1887–1897. (doi:10.1242/jeb.00372.)
- Stephens, D. W. & Krebs, J. R. 1986 *Foraging theory*. Princeton, NY: Princeton University Press.
- Tomkovich, P. S. 2001 A new subspecies of red knot *Calidris canutus* from the New Siberian Islands. *Bull. Brit. Ornithol. Club* **121**, 257–263.
- Van der Meer, J. & Piersma, T. 1994 Physiologically inspired regression models for estimating and predicting nutrient stores and their composition in birds. *Physiol. Zool.* **67**, 305–329.
- Van Gils, J. & Piersma, T. 1999 Day- and nighttime movements of radiomarked knots, *Calidris canutus*, staging in the western Wadden Sea in July–August 1995. *Wader Study Group Bull.* **89**, 36–44.
- Van Gils, J., Piersma, T., Dekinga, A. & Spaans, B. 2000 Distributional ecology of individually radio-marked Knots *Calidris canutus* in the western Dutch Wadden Sea in August–October 1999. *Limosa* **73**, 29–34.
- Van Gils, J. A., Piersma, T., Dekinga, A. & Dietz, M. W. 2003 Cost-benefit analysis of mollusc-eating in a shorebird. II. Optimizing gizzard size in the face of seasonal demands. *J. Exp. Biol.* **206**, 3369–3380. (doi:10.1242/jeb.00546.)
- Van Gils, J. A., De Rooij, S. R., Van Belle, J., Van der Meer, J., Dekinga, A., Piersma, T. & Drent, R. 2005a Digestive bottleneck affects foraging decisions in red knots (*Calidris canutus*). I. Prey choice. *J. Anim. Ecol.* **74**, 105–119. (doi:10.1111/j.1365-2656.2004.00903.x.)
- Van Gils, J. A., Dekinga, A., Spaans, B., Vahl, W. K. & Piersma, T. 2005b Digestive bottleneck affects foraging decisions in red knots (*Calidris canutus*). II. Patch choice and length of working day. *J. Anim. Ecol.* **74**, 120–130. (doi:10.1111/j.1365-2656.2004.00904.x.)
- Weiner, J. 1992 Physiological limits to sustainable energy budgets in birds and mammals: ecological implications. *Trends Ecol. Evol.* **7**, 384–388. (doi:10.1016/0169-5347(92)90009-Z.)
- Wetlands International 2002 *Waterbird population estimates*. 3rd edn. Wetland International Global Series 12, Wageningen, The Netherlands.
- Wiersma, P. & Piersma, T. 1994 Effects of microhabitat, flocking, climate and migratory goal on energy expenditure in the annual cycle of red knots. *Condor* **96**, 257–279.
- Wikelski, M., Tarlow, E. M., Raim, A., Diehl, R. H., Larkin, R. P. & Visser, G. H. 2003 Cost of migration in free-flying songbirds. *Nature* **423**, 704. (doi:10.1038/423704a.)
- Wilson, B. R. 1969 Survival and reproduction of the mussel *Xenostrobus securis* (Lamarck) (Mollusca; Bivalvia; Mytilidae) in a western Australian estuary. Pt. II. Reproduction, growth and longevity. *J. Nat. Hist.* **3**, 93–120.
- Wilson, B. R. & Hodgkin, E. P. 1967 A comparative account of the reproductive cycles of five species of marine mussels (Bivalvia: Mytilidae) in the vicinity of Fremantle, western Australia. *Aust. J. Mar. Fresh. Res.* **18**, 175–203.
- Yearsley, J., Hastings, I. M., Gordon, I. J., Kyriazakis, I. & Illius, A. W. 2002 A lifetime perspective on foraging and mortality. *J. Theor. Biol.* **215**, 385–397. (doi:10.1006/jtbi.2002.2529.)
- Zwarts, L. 1991 Seasonal variation in body weight of the bivalves *Macoma balthica*, *Scrobicularia plana*, *Mya arenaria* and *Cerastoderma edule* in the Dutch Wadden Sea. *Neth. J. Sea Res.* **28**, 231–245. (doi:10.1016/0077-7579(91)90021-R.)
- Zwarts, L. & Wanink, J. H. 1993 How the food supply harvestable by waders in the Wadden Sea depends on the variation in energy density, body weight, biomass, burying depth and behaviour of tidal-flat invertebrates. *Neth. J. Sea Res.* **31**, 441–476. (doi:10.1016/0077-7579(93)90059-2.)
- Zwarts, L., Blomert, A.-M. & Hupkes, R. 1990 Increase of feeding time in waders preparing for spring migration from the Banc d'Arguin, Mauritania. *Ardea* **78**, 237–256.

The electronic supplementary material is available at <http://dx.doi.org/10.1098/rspb.2005.3245> or via <http://www.journals.royalsoc.ac.uk>.

As this paper exceeds the maximum length normally permitted, the authors have agreed to contribute to production costs.